

Exploring MIP Sensor of Basal Stem Rot (BSR) Disease in Palm Oil Plantation

A.H. Abdullah¹, A.Y.Md. Shakaff¹, A.H. Adom¹, M.N. Ahmad¹, A. Zakaria¹, S.A. Ghani¹, N.M. Samsudin¹, F.S.A. Saad¹, L.M. Kamarudin¹, N.H. Hamid², I.A. Seman³

¹Center of Excellence Advanced Sensor Technology, UniMAP, 01000 Kangar, Perlis, Malaysia
e-mail:abu.hassan@unimap.edu.my

²Felda Agriculture Services SdnBhd (FASSB), Jalan Gurney Satu, 54000 Kuala Lumpur, Malaysia

³Malaysian Palm Oil Board (MPOB), MPOB, Bandar BaruBangi, 43000 Kajang, Selangor, Malaysia

Abstract:

In Malaysia, the production of palm oil is hampered by the infection of Basal Stem Rot (BSR) disease. Unfortunately, the existing BSR detection techniques are complex, time-consuming and still not fully developed. So, this research proposes an investigation to identify the volatile compounds (biomarkers) of the infected oil palm trees. The biomarkers will be proposed to develop specific Molecularly Imprinted Polymer (MIP) sensors. The sensor will be used for the development of 'Application Specific Electronic Nose' (ASEN). The samples were taken from healthy and infected BSR disease tree. Fourier Transform Infrared (FTIR) and Headspace Gas Chromatography Mass Spectrometry (GC-MS) analysis is used to identify the biomarkers. The results show that the technique has successfully detected and identified the volatile compounds *biomarkers* for BSR disease of infected oil palm tree.

Key words: BSR disease, FTIR, GC-MS, MIP and e-nose.

Introduction

Oil palm is a tropical perennial species which mainly used in foods, detergents and lubricants. Now the demand is soar because it is also being used for biofuel. Malaysia has 4.2 million hectares of palm oil that enables it to exceed natural rubber as the main cash crop [1]. The crop is susceptible to diseases, commonly fungus, pests and pathogens where Basal Stem Rot (BSR) is the main disease in Malaysia and Indonesia [2]. The *Ganoderma boninense* fungus is the sole causal agent of the disease in Malaysia [3]. The disease infection occurs when health roots contact with diseases tissues in the soil, wounded tissues or dead roots [4]. Then the fungus destroys the internal palm tissues, which eventually leads to the death of the tree [5]. It is estimated that if 2.5 per cent of the total acreage of oil palm plantation in Malaysia is affected by the disease, the industry would lose about RM80 million each year [6].

Currently, numbers of research works have been reported on the detection of *Ganoderma boninense* fungus and the BSR disease. The most common technique is the field symptom known as 'skirting' where some of the mature leave start to wilt, malnutrition and going down. The disease can also be detected from the rotten palm trunk base and the presence of

small white button fruiting bodies (basidiomata) [7]. Other techniques include morphological characteristics, pathogenicity test, Polymerase Chain Reaction (PCR), Enzyme-Linked Immunosorbent Assay (ELISA), tomography and hyper-spectral reflectance data.

Unfortunately most of these techniques require complex procedure and still not fully developed. The techniques can only detect BSR by the presence of the fruiting bodies on the palm trunk base (see Fig. 1). At this stage the disease cannot be cured and the tree had already dead. Subsequently, the disease might spread to next adjacent trees. In addition, those techniques are also unable to provide real time results. The availability of speedy and accurate diagnostic techniques, for detecting *Ganoderma boninense* in oil palm at an early stage of infection, would benefit the industry.

The research is exploring the possibilities of identifying the BSR volatile compounds main chemical components (biomarkers) by using Fourier Transform Infrared (FTIR) and Headspace Gas Chromatography Mass Spectrometry (GC-MS). Specific chemical standard will be used to verify the biomarkers. The *Ganoderma*-selective medium (GSM) tissue culture will be used to validate the GC-MS analysis result. The biomarkers also will be

further quantified to the degree of the *Ganoderma boninense* infection. The biomarkers will be used to develop specific Molecularly Imprinted Polymer (MIP) sensors. The imprinting molecule is used as a template to form a receptor site with complementary binding features with regard to the molecules of the volatile compounds. The MIP substance contains mixed chemical compounds that detect the biomarkers. The MIP is coated on Quartz Crystal Microbalance (QCM) where conducting polymer being used for binding. The sensor will be tested and characterise to verify the functionality.



Fig.1: The *Ganoderma boninense* basidiomata.

The MIP sensors are used to develop an 'Application Specific Electronic Nose' (ASEN). The instrument consists of sensors, sensing chamber and microcontroller with embedded Artificial Neural Network (ANN) pattern recognition. The instrument will provide on-site early detection of the BSR disease in oil palm plantation.

Methodology

Sample of skin, soil, fruiting body and air around the stump are taken from healthy and infected BSR disease tree. The samples are collected from the Federal Land Development Authority (FELDA) palm oil Research Station in Malaysia. The tree is being identified by BSR disease expert (FELDA staff with 20 years working experience in the oil palm plantation) by using field symptoms technique. The tree is being identified by *Ganoderma* expert (plant pathologist) and verify by using ELISA-PAB technique.

The Fourier Transform Infrared (FTIR) Spectroscopy is used to identify the *Ganoderma boninense* functional group. The FTIR spectrum has been obtained by using a Perkin Elmer 2000 Series Instrument (Perkin – Elmer Cetus Instruments, Norwalk CT). The spectrum resolution is set at 4 cm^{-1} and the scanning range was selected from 600 to 4000

cm^{-1} . The result of the analysis will be used to strategies the GC-MS analysis.


The GC-MS analysis is performed offline by using GC-MS Clarus 600T (Perkin Elmer, Massachusetts, USA) for the infected and healthy trunk sample. Five gram of each sample was added directly into 22ml headspace vial and seal with PTFE septa. The samples were kept in the glass jar for 6 hours to allow volatiles to concentrate before each measurement. The sample was scan of total ion current strength (TIC) from m/z 40-500 with incubation triplicate in Turbo Matrix Headspace Sampler for 60 minutes at 80°C . Then it is analysed using Clarus 680 GC for separation and Clarus 600T for mass analysis. The simplest data time is basically a chromatographic output representing a summation of the signal output from the mass spectrometer analyser which is a measurement of the TIC versus strength of all the ions produced by the mass spectrometer at a given time. The headspace compound identification is done by looking at the retention time and comparing with the known library standard and search hits.

Results and discussion


The FTIR spectrum result (see Fig. 2) and Table 1) shows summaries of the *Ganoderma* FTIR spectrum functional group analysis. The percentage transmittance of the absorption analysis has been identified and the *Ganoderma* functional groups were carbohydrate, hemicellulose and lignocellulose.

Table 1: Functional Group of *Ganoderma*

W.length (cm^{-1})	Functional Group	Organic Compound
3846.96	–OH, (Hydroxyl)	9(11),22-trien-3b-ol, 2b,3a,9a-trihydroxyergosta-7,22-diene
3003.63	CH_3 (Methyl) or $-\text{CH}_2$ (Methylene)	All the compound in table 4
2263.85	CN (Cyanide)	Little toxic odour
1567.63	Aromatic ring	All the compound in table 4
1190.63	C-O-C	ergosta-7,22-dien-3b-yl linoleate, ergosta-7,22-dien-3b-yl palmitate
697. 63	Si-O (Silicone)	Small quantity for odour binding



The GC-MS analysis result on the headspace volatile of the infected tree and non-infected tree trunk is able to identify the *Ganoderma boninense* biomarkers (see Fig. 3(a), (b) and 3(c)). Two peaks of the result were identified as the volatile biomarkers of the infected BSR disease. MS search on these two peaks found that Benzene,1-Methyl-2-(2-Propenyl)- and Cyclohexasiloxane, dodecamethyl-. Calibration with standard chemicals will further ensure the integrity of these findings.



Conclusion

This research is successfully identifying the volatile compounds component (biomarkers) for *Ganoderma boninense* fungus of the BSR infected oil palm trees. Fourier Transform Infrared (FTIR) and Headspace Gas Chromatography Mass Spectrometry (GCMS) are used to identify the *biomarkers*. Specific chemical standard will be used to verify the biomarkers. The finding will be used to develop

specific MIP sensors for 'Application Specific Electronic Nose' (ASEN) e-nose for detecting the degree of BSR infection. The results demonstrate that the technique has successfully detected and identified the volatile compound biomarkers for BSR disease of oil palm tree at high accuracy rate. This will enable preventive action to be taken to prolong the oil palm tree life cycle as well as preventing the disease from spreading to the whole plantation.

Acknowledgements

This project is partly funded by the Malaysian Ministry of Higher Education under the Fundamental Research Grant Scheme (FRGS9003-00250). A.H. Abdullah gratefully acknowledges the leave granted by Universiti Malaysia Perlis (UniMAP) to allow him to pursue his Ph.D.

References

- [1] A.C. Soh, C.K. Wong, Y.W. Ho, and C.W. Choong, Oil Palm J. Vollmann, I. Rajcan (eds.), Oil Crops, *Handbook of Plant Breeding 4*, (2010): doi: 10.1007/978-0-387-77594-4-11
- [2] Azahar T. M., Mustapha, J.C., Mazliham, S. and Boursier, Temporal Analysis of Basal Stem Rot Disease in Oil Palm Plantations: An Analysis on Peat Soil, *Int. Journal of Engineering & Technology IJET-IJENS. Vol: 11 No: 03, (2011)*.
- [3] A.C.Soh, C.K. Wong, Y.W.Ho, and C.W.Choong, "Oil Crops, *Handbook of Plant Breeding 4*", Oil Palm J. Vollmann, Springer Science Business Media, LLC, (2009).
- [4] PD Turner, Oil Palm and Ganoderma III: Treatment and Control in Established Plantings, *The Planter*, 41:279-282, (1965).
- [5] M. A. Markoma, A. Y. Md. Shakaff, A. H. Adom, M. N. Ahmad, W. Hidayat, A. H. Abdullah, and N. Ahmad Fikri, *Computers and Electronics in Agriculture* 66, 140 (2009).
- [6] H Singh, MN Ahmad, MZ Daud, F Khairul, AYM Shakaff and S Hamzah, *Electronic Nose Application In Ganoderma Detection?*, 1st. Research Seminar, KUKUM, Malaysia, (2006).
- [7] CD Lelong, JM Roger, S Brégand, F Dubertret, Evaluation of Oil-Palm Fungal Disease Infestation with Canopy Hyperspectral Reflectance Data, *Sensors*, 10, 734-747, (2010).